

Oxalic Acid in Varroa Management

**Randy Oliver
Revised 14 Jan 2016**

(see notes under the slides)

Oxalic acid for the control of varroosis in honey bee colonies – a review¹

Eva RADEMACHER*, Marika HARZ

Free University of Berlin, Dept. of Biology/Chemistry/Pharmacy, Neurobiology, Königin-Luise-Str. 28–30,
14195 Berlin, Germany

Received 17 February 2005 – revised 17 June 2005 – accepted 26 July 2005

Abstract – The review summarizes research results on the use of oxalic acid as an acaricide in honey bee colonies. Three different treatment techniques (i.e. trickling, evaporation and spraying) have been developed for the application of oxalic acid. Detailed information is given on the efficacy against *Varroa destructor*, tolerability by *Apis mellifera*, protective procedures for the user, residue situation and consumer safety, as well as recommendations for use.

Oxalic has a long history in Europe

Why Oxalic Acid?




**Acids are much more
toxic to mites than to bees**

Oxalic is the strongest organic acid

pH of Organic Acids – Sorted by increasing pH

Formula	Name	Alternative name	1 mM	10 mM	100 mM
C ₂ H ₂ O ₄	oxalic acid	ethanedioic acid	3.00	2.09	1.31
C ₄ H ₄ O ₄	maleic acid	toxic acid	3.05	2.21	1.54
C ₄ H ₆ O ₈	dihydroxytartaric acid		3.04	2.20	1.54
C ₆ H ₆ O ₆	cis-aconitic acid		3.05	2.21	1.55
C ₄ H ₄ O ₅	oxaloacetic acid	oxalacetic acid	3.05	2.29	1.68
C ₅ H ₆ O ₅	oxoglutaric acid	ketoglutaric acid	3.09	2.36	1.77
C ₃ H ₄ O ₃	pyruvic acid	acetylformic acid	3.11	2.38	1.79
C ₄ H ₆ O ₃	2-oxobutanoic acid	alpha-ketobutyric acid	3.11	2.38	1.79
C ₆ H ₆ O ₆	trans-aconitic acid		3.14	2.48	1.92
C ₃ H ₄ O ₄	malonic acid	propanedioic acid	3.17	2.50	1.94
C ₈ H ₆ O ₄	phthalic acid	ortho-phthalic acid	3.20	2.55	2.00
C ₄ H ₆ O ₆	L-tartaric acid	racemic acid	3.18	2.55	2.01
C ₇ H ₆ O ₄	gentisic acid	"2,5-dihydroxybenzoic"	3.21	2.56	2.01
C ₇ H ₆ O ₃	salicylic acid	2-hydroxybenzoic acid	3.21	2.57	2.02
C ₄ H ₄ O ₄	fumaric acid	allomaleic acid	3.19	2.57	2.03
C ₄ H ₆ O ₆	DL-tartaric acid	racemic acid	3.19	2.58	2.03
C ₄ H ₆ O ₄	methylmalonic acid		3.23	2.60	2.05
C ₅ H ₆ O ₄	mesaconic acid	dimethylfumaric acid	3.23	2.61	2.06
C ₄ H ₆ O ₆	meso-tartaric acid	racemic acid	3.24	2.62	2.08
C ₆ H ₈ O ₇	citric acid		3.24	2.62	2.08
C ₂ H ₂ O ₃	glyoxilic acid	oxoacetic acid	3.27	2.65	2.11
C ₆ H ₈ O ₇	isocitric acid		3.28	2.69	2.16
C ₄ H ₆ O ₅	malic acid		3.33	2.74	2.21
C ₈ H ₈ O ₃	mandelic acid	phenylglycolic acid	3.34	2.75	2.22
C ₈ H ₆ O ₄	isophthalic acid	meta-phthalic acid	3.33	2.76	2.24
C ₈ H ₆ O ₄	terephthalic acid	para-phthalic acid	3.36	2.79	2.26
C ₃ H ₆ O ₄	glyceric acid		3.38	2.80	2.27
C ₄ H ₆ O ₃	acetoacetic acid	diacetic acid	3.40	2.83	2.30
C ₄ H ₈ O ₃	hydroxybutanoic acid		3.43	2.86	2.33
CH ₂ O ₂	formic acid	methanoic acid	3.47	2.91	2.38
C ₂ H ₄ O ₃	glycolic acid	hydroxyacetic acid	3.50	2.94	2.42
C ₅ H ₆ O ₄	itaconic acid	methylenesuccinic acid	3.50	2.95	2.43
C ₃ H ₆ O ₃	lactic acid	milk acid	3.51	2.96	2.44
C ₈ H ₈ O ₂	toluic acid	ortho-toluic acid	3.53	2.98	2.46
C ₆ H ₈ O ₆	ascorbic acid	vitamin C	3.59	3.04	2.53
C ₅ H ₈ O ₄	methylsuccinic acid		3.62	3.08	2.57
C ₇ H ₆ O ₂	benzoic acid		3.66	3.12	2.60
C ₈ H ₈ O ₂	o-toluic acid	ortho-toluic acid	3.65	3.13	2.61



<http://www.varroamilbe.ch>

There are speculative hypotheses as to why acids kill varroa, but no definitive study.

Beekeeper Gerhard Bruning suspects that OA crystals are absorbed through varroa's sticky tarsal pads.

Safety to Humans

Oxalis






Table I: Oxalic acid content in selected vegetables. These are ballpark numbers, actual content depends on many variables. See note at the end of the article.

Vegetable	Oxalic acid (g/100 g)
Asparagus	0.13
Broccoli	0.19
Lettuce	0.33
Brussels sprouts	0.36
Collards	0.45
Beet leaves	0.61
Spinach	0.97 (0.32-1.26)
Purslane	1.31
Parsley	1.70
Rhubarb	0.3-1.5
leaf	0.59 - 0.72
stalk	0.39 - 0.54

Typical treatment=
1 serving
per hive



Purchasing Oxalic Acid



Sold as
oxalic acid
dihydrate

“Wood Bleach”



Serving the industry for more than 30 years

Brushy Mountain Bee Farm

BEST QUALITY | BEST SERVICE | BEST SUPPORT

[Home](#) [Quick Order](#) [Request Catalog](#) [Bee Educated](#) [Account Login](#)

Online Store

[Beekeeping Supplies](#)

[Candle Making Supplies](#)

[Soap and Skin Care Supplies](#)

[Mead and Wine Supplies](#)

[Gifts](#)

[Monthly Specials](#)

[Bargain Aisle - Discontinued and Scratch/Dent](#)

[Featured Products](#)



[New for 2015](#)

[Cyber Monday](#)

Free Shipping on Most Orders Over \$150*

Free Shipping does not apply to international orders, back ordered items shipped separately, glass jars, honey or syrup. LTL shipments will be sent out at no charge but any accessorial fees will be billed to customer's account. Other restrictions may apply.

Search Results for : "oxalic"

Item #	Image	Description
727		Oxalic Acid Shp. Wt. 1 lb.
727K		Kit for Oxalic Acid Ship wt. 2 lbs. In Stock
727OAV		Varrocleaner Ship wt. 3 lbs. In Stock

This is the only registered and legal oxalic product!

Mixing Oxalic Syrup

American Weigh Scale

0.1 G

by American Weigh Scale

\$9.89 ~~\$20.95~~  Prime

Get it by **Monday, Nov 30**


More Buying Choices

\$6.00 new (8 offers)



Need to
weigh

HARD WATER LEVELS THROUGHOUT THE U.S.



LEGEND

- 0 - 3.5 grains per gallon – Soft
- 3.5 - 7.0 grains per gallon – Moderately Hard
- 7.0 - 10.5 grains per gallon – Hard
- over 10.5 grains per gallon – Very Hard

Due to the nature of the water bed sub-soil structure, water hardness may vary from one source to another within a general area.

Don't use hard water



ScientificBeekeeping.com

Beekeeping Through the Eyes of a Biologist

[Home](#) [Contact Me](#)

Oxalic Acid Treatment Table

[Save](#) [PDF](#) [Email](#) [Print](#)

Be sure to read "[Oxalic Acid – Questions and Answers](#)" and "[The Learning Curve – Part 3](#)" before using oxalic acid. It is critical to apply it correctly, or you risk seriously harming your bees!"

Important Note: the following proportions refer to common oxalic acid dihydrate (wood bleach). If you manage to get your hands on pure laboratory oxalic acid, you must reduce the amount of acid to only 7/10ths of that of the dihydrate !!!!

Also note that if you use hard water, some of the oxalic acid will precipitate out as calcium oxalate, and thus reduce the efficacy (you'll easily see the white precipitate if this is the case).

Oxalic strength →	"Hot" 4.2% w:v	"Medium" 3.2% w:v	"Weak" 2.5% w:v	Notes
OA Crystals	1	0.75	0.6	Oxalic crystals must be measured by weight. Sugar and water are about the same by weight or volume (1 pint of either granulated sugar or water weight 1 lb)
Sucrose	10	10	10	
Dist. Water	10	10	10	
OA Crystals	60g	45g	35g	Makes 1 liter;

[Articles By Publication Date](#)[Bee Behavior and Biology](#)[Varroa Management](#)[Bee Nutrition](#)[Colony Health – Diseases, Viruses, CCD](#)[Pesticide Issues](#)[Nosema ceranae](#)[Almond Pollination](#)[Miscellaneous articles](#)[Research Updates](#)[Beginner's Pages](#)[Please Share](#)

Use the exact dose!

<i>Oxalic strength→</i>	<i>“Hot” 4.2% w:v</i>	<i>“Medium” 3.2% w:v</i>	<i>“Weak” 2.5% w:v</i>	Notes
OA crystals	1	0.75	0.6	Oxalic crystals must be measured by weight. Sugar and water are about the same by weight or volume (1 pint of either granulated sugar or water weigh 1 lb)
Sucrose	10	10	10	
Dist. Water	10	10	10	
OA crystals	60g	45g	35g	
Sucrose	600g	600g	600g	Makes 1 liter Treats about 20 colonies
Dist. water	600ml	600ml	600ml	
OA crystals	100g	75g	60g	
Sucrose	1 kg	1 kg	1 kg	Makes 1700ml Treats about 33 colonies
Dist. water	1 liter	1 liter	1 liter	
OA crystals	232g	174g	139g	
Sucrose	5 lb	5 lb	5 lb	Makes 1+ gallon Treats about 75 colonies
Dist. water	2.5 qt	2.5 qt	2.5 qt	
OA crystals	1112g (2lb 7oz)	834g (1lb 13.4oz)	667g (1lb 7.5oz)	
Sucrose	25 lb	25 lb	25 lb	Makes 5 gallons Treats about 375 colonies
Dist. water	3 gal	3 gal	3 gal	

Oxalic acid crystals dissolve more readily in hot water than in sugar solution.

Tip: dissolve the oxalic crystals in the indicated amount of hot (150°F) water before adding the sugar.

After the oxalic crystals are *fully dissolved*, *only then* stir in the sugar.

Storage

Store in the 'fridge



Safety



Protect your eyes



**Tastes like strong
lemonade**

DIY Alkaline Water

homeAIDmom.blogspot.com

Carry baking
soda in water to
neutralize



Application



**Must be applied directly
to bees' bodies.**

~5 mL per "seam" of
bees



1 tsp = 5 mL





Dribble, not spray

**Calibrate pump
output**





Hit both boxes

Tips:

Fill the garden sprayer only about $\frac{1}{4}$ full of solution. This leaves a large air space, which minimizes the fluctuation in pressure.

After you've dribbled a yard of hives, measure how much syrup you've applied in total, and divide by the number of hives. This will tell you if you're applying the correct amount.


Timing of Treatment



Oxalic won't kill mites in the brood.

Kill rate of phoretic mites					
90	← you can change this value				
Percent of full broodnest	Estimated percentage of mites in the brood	Estimated percentage of mites phoretic	Kill rate of phoretic mites	Percent kill of total mite population	
100	70%	30%	90%	27%	
90	63%	37%	90%	33%	
80	56%	44%	90%	40%	
70	49%	51%	90%	46%	
60	42%	58%	90%	52%	
50	35%	65%	90%	59%	
40	28%	72%	90%	65%	
30	21%	79%	90%	71%	
20	14%	86%	90%	77%	
10	7%	93%	90%	84%	
0	0%	100%	90%	90%	


Note the difference in efficacy, dependent upon how much brood is present.



Oxalic gives poor efficacy if there is much brood present, especially if drone brood is present.

Age Class Distribution of Workers Over the Year Manitoba, Shed Wintered

© Randy Oliver 2015



Best treatment windows



Fall treatment



Oxalic acid is, by far, most effective when colonies are broodless.





**We use our fall oxalic dribble
as a last check on our
colonies before winter.**



**Oxalic drops mites
for about 4 days.**

Benefit against nosema

Accepted Manuscript


Effect of oxalic acid on *Nosema ceranae* infection

Antonio Nanetti, Cristina Rodriguez-García, Aránzazu Meana, Raquel Martín-Hernández, Mariano Higes

PII: S0034-5288(15)30033-3

DOI: [doi: 10.1016/j.rvsc.2015.08.003](https://doi.org/10.1016/j.rvsc.2015.08.003)






Nosema infection after fall dribble of weak OA, 50mL/hive

Summer Treatment

- *Can be used on severely mite-stressed colonies to buy time.*
- *Must be repeated at weekly intervals.*

**Colony about to
collapse**






Formic or
thymol may be
too strong a
treatment



**Summer
treatment—
3 weekly
applications.**

**Around 50% mite
reduction.**

Induced Brood Break



Understand the timing!




Beekeepers in Italy create an induced brood break during late summer by temporarily caging the queen

Creating an induced brood break for oxalic treatment by caging the queen for two weeks.

Cage Queen

Release Queen

Treat with oxalic





Cage the queen for 12 days minimum. Then release her.
Treat 4-5 days later.



Treat walkaway splits at 20 days.

Combine OA with requeening



Kill the old queen,
insert a queen
cell,
treat with OA 19
days later.



Cage the old queen for 2 weeks,
then remove her and introduce a new queen,
treat with OA 5 days later.

Treatment of Nucs or Packages


A no brainer




Treat package bees or swarms shortly after installation for a "clean start"

Treatment window for nucs

The Biology Behind the Method




Treatment Effect upon Normalized Mite Infestation Rates over 3 Months




Timeline, Trial 2


Oxalic dribble of nucs



Trial 2: Mite Infestation Rate in Nucs 53 Days Post Treatment



Trial 2: Colony Strength 53 Days Post Treatment





Doesn't appear to harm queens


Graft #	Queen(s)	No. cells	Day	Graft date	Sealed	Incubator	Nucs by	Early ripe	Typ mate	Typ eggs	Check/oxalic	Nuc yard(s)
			Fri	3/9	3/13	3/16	3/18	3/19	3/28	3/31	4/6	
			Sat	3/10	3/14	3/17	3/19	3/20	3/29	4/1	4/7	
			Sun	3/11	3/15	3/18	3/20	3/21	3/30	4/2	4/8	
			Mon	3/12	3/16	3/19	3/21	3/22	3/31	4/3	4/9	
			Tue	3/13	3/17	3/20	3/22	3/23	4/1	4/4	4/10	
			Wed	3/14	3/18	3/21	3/23	3/24	4/2	4/5	4/11	
			Th	3/15	3/19	3/22	3/24	3/25	4/3	4/6	4/12	

Create a spreadsheet to keep track of dates

Vaporization (Sublimation)

A Magic Wand?





Useful where winter comes on suddenly.



Simple Varroa vaporizer.



**Vaporized oxalic is dangerous!
Wear a respirator.**

Other issues:
Cooked bees
Burnt wood or plastic





Sublimation may be easier on the bees.

Recently published

Journal of Apicultural Research, 2015
<http://dx.doi.org/10.1080/00218839.2015.1106777>



ORIGINAL RESEARCH ARTICLE

Towards integrated control of varroa: 2) comparing application methods and doses of oxalic acid on the mortality of phoretic *Varroa destructor* mites and their honey bee hosts

Hasan Al Toufailia*, Luciano Scandian and Francis L W Ratnieks



Dribble vs. Sublimation

Dribble		Sublimation	
Pros:	High efficacy	Pros:	Perhaps higher efficacy
	Very safe to apply		No opening of the hive
	Quick		Can do in freezing weather
	Little equip needed		Perhaps gentler to the bees
			No syrup mixing
Cons:	Requires opening hive	Cons:	Vapor fog is hazardous
	May be problematic in freezing weather		Requires specialized vaporizer and energy source
	Easier with helper		Problems with hot tip

Instructions: type values into for the two yellow cells and hit enter


Percent of full broodnest present 0 % A full broodnest with substant

there may be only 50% or less of a full broodnest present. Adjust this value to

Expected weekly kill of the phoretic mites by the treatment 90 % fil

See [f](#)

Effect of a single treatment, no brood



© Randy Oliver 2015

Instructions: type values into for the two yellow cells and hit enter


Percent of full broodnest present 100 % A full broodnest with substantial

there may be only 50% or less of a full broodnest present. Adjust this value to

Expected weekly kill of the phoretic mites by the treatment 90 % fill

See No

Effect of a single treatment, full broodnest



© Randy Oliver 2015

Instructions: type values into for the two yellow cells and hit enter


Percent of full broodnest present 100 % A full broodnest with substantial

there may be only 50% or less of a full broodnest present. Adjust this value to

Expected weekly kill of the phoretic mites by the treatment 90 % fill

See No

Effect of treatment once every three weeks



© Randy Oliver 2015

Instructions: type values into for the two yellow cells and hit enter


Percent of full broodnest present 100 % A full broodnest with substantial

there may be only 50% or less of a full broodnest present. Adjust this value to yc

Expected weekly kill of the phoretic mites by the treatment 90 % fill in

See Not

Effect of weekly treatment



© Randy Oliver 2015

**Another application
method**

A new formulation of oxalic acid for *Varroa destructor* control applied in *Apis mellifera* colonies in the presence of brood

Matías MAGGI^{1,2}, Elian TOURN^{3,4,5}, Pedro NEGRI^{1,2}, Nicolás SZAWARSKI¹,
Alfredo MARCONI^{3,4,5}, Liliana GALLEZ⁶, Sandra MEDICI^{1,2}, Sergio RUFFINENGO⁷,
Constanza BRADESCO¹, Leonardo De FEUDIS¹, Silvina QUINTANA⁸, Diana SAMMATARO⁹,
Martin EGUARAS^{1,2}




Oxalic/glycerine on cardboard strips.

A Critical Closing Thought

“The only way to halt the development of resistance to a certain product is by interrupting its use in the control strategy.”

Lodesani (2009) Limits of chemotherapy in beekeeping: development of resistance and the problem of residues.

Practice some sort of rotation of treatments





Happy beekeeping!
ScientificBeekeeping.com